

## Reference List

- AGS. (2001). Guideline for the prevention of falls in older persons. American Geriatrics Society, British Geriatrics Society, and American Academy of Orthopaedic Surgeons Panel on Falls Prevention. *J Am Geriatr Soc*, 49(5), 664-672.
- Baker, L. (1992). Developing a safety plan that works for patients and nurses. *Rehabil Nurs*, 17(5), 264-266.
- Balducci, L. (2007). Aging, frailty, and chemotherapy. *Cancer Control*, 14(1), 7-12.
- Barnes, D. E., Palmer, R. M., Krevsevic, D. M., Fortinsky, R. H., Kowal, J., Chren, M. M., & Landefeld, C. S. Acute care for elders units produced shorter hospital stays at lower cost while maintaining patient' functional status. *Health*, 31(6), 1227-36.
- Berg, K. O., Wood-Dauphinee, S. L., Williams, J. I., & Maki, B. (1992). Measuring balance in the elderly: validation of an instrument. *Can J Public Health*, 83 Suppl 2, S7-11.
- Bernabeu-Mora, R., Medina-Mirapeix, F., Llamazares-Herran, E., Garcia-Guillmon, G., Gimenez-Gimenez, L., Sanchez-Neito, J. (2015). The short physical performance battery is a discriminative tool for identifying patients with COPD at risk of disability. *International Journal of COPD*, 10,2619-2626.
- CDC. (2006). National Center for Injury Prevention and Control: Web-based Injury Statistics Query and Reporting System (WISQARS). [www.cdc.gov/ncipc/wisqars](http://www.cdc.gov/ncipc/wisqars).
- Chen, H., Cantor, A., Meyer, J., Beth Corcoran, M., Grendys, E., Cavanaugh, D., . . . Extermann, M. (2003). Can older cancer patients tolerate chemotherapy? A prospective pilot study. *Cancer*, 97(4), 1107-1114.
- Cumming, R. G. (1998). Epidemiology of medication-related falls and fractures in the elderly. *Drugs Aging*, 12(1), 43-53.
- Dolinis, J., Harrison, J. E., & Andrews, G. R. (1997). Factors associated with falling in older Adelaide residents. *Aust N Z J Public Health*, 21(5), 462-468.
- Fairhill, N., Langron, C., Sherrington, C., Lord, S., Kurrle, S., Lockwood, K., Monaghan, N., Aggar, C., Gill, L., & Cameron, I., (2011). Treating frailty-a practical guide. *BMC Medicine*, 9(83).
- Fried, L. P., Tangen, C. M., Walston, J., Newman, A. B., Hirsch, C., Gottdiener, J., . . . McBurnie, M. A. (2001). Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci*, 56(3), M146-156.
- Given, B., Given, C., Azzouz, F., & Stommel, M. (2001). Physical functioning of elderly cancer patients prior to diagnosis and following initial treatment. *Nurs Res*, 50(4), 222-232.
- Guralnik, J., Simonsick, E.M., Ferrucci, L., Glynn, R., Berkman, L.F., Blazer, D., Scherr, P.A., & Wallace, R. B. (1994). A short physical performance battery assessing lower extremity function: Association with self-reported disability and prediction of mortality and nursing home placement. *Journal of Gerontology*, 49(2), M85-M94.
- Handforth, C., Clegg, A., Young, C., Simpkins, S., Seymour, M. T., Selby, P. J., & Young, J. (2015). The prevalence and outcomes of frailty in older cancer patients: a systematic review. *Ann Oncol*, 26(6), 1091-1101. doi: 10.1093/annonc/mdu540

- Hausdorff, J. M., Rios, D. A., & Edelberg, H. K. (2001). Gait variability and fall risk in community-living older adults: a 1-year prospective study. *Arch Phys Med Rehabil*, 82(8), 1050-1056.
- Holley, S. (2002). A look at the problem of falls among people with cancer. *Clin J Oncol Nurs*, 6(4), 193-197.
- Kario, K., Tobin, J. N., Wolfson, L. I., Whipple, R., Derby, C. A., Singh, D., . . . Wassertheil-Smoller, S. (2001). Lower standing systolic blood pressure as a predictor of falls in the elderly: a community-based prospective study. *J Am Coll Cardiol*, 38(1), 246-252.
- Katz, S., Downs, T. D., Cash, H. R., & Grotz, R. C. (1970). Progress in development of the index of ADL. *Gerontologist*, 10(1), 20-30.
- Kellogg International Work Group. (1987). A report of the Kellogg International Work Group on the Prevention of Falls by the Elderly. *Dan Med Bull*, 34 Suppl 4, 1-24.
- Kelsey, J. L., Procter-Gray, E., Nguyen, U. S., Li, W., Kiel, D. P., & Hannan, M. T. (2010). Footwear and Falls in the Home Among Older Individuals in the MOBILIZE Boston Study. *Footwear Sci*, 2(3), 123-129. doi: 10.1080/19424280.2010.491074
- Kim, Y. H., Kim, K. I., Paik, N. J., Kim, K. W., Jang, H. C., & Lim, J. Y. (2016). Muscle strength: A better index of low physical performance than muscle mass in older adults. *Geriatr Gerontol Int*, 16(5), 577-585. doi: 10.1111/ggi.12514
- Koski, K., Luukinen, H., Laippala, P., & Kivela, S. L. (1996). Physiological factors and medications as predictors of injurious falls by elderly people: a prospective population-based study. *Age Ageing*, 25(1), 29-38.
- Kutner, N. G., Zhang, R., Huang, Y., & Painter, P. (2015). Gait Speed and Mortality, Hospitalization, and Functional Status Change Among Hemodialysis Patients: A US Renal Data System Special Study. *Am J Kidney Dis*. doi: 10.1053/j.ajkd.2015.01.024
- Lau, E. M., Woo, J., & Lam, D. (1991). Neuromuscular impairment: a major cause of non-syncopal falls in elderly Chinese. *Public Health*, 105(5), 369-372.
- Lawton, M. , & Brody, E. M. (1969). Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist*, 9(3), 179-186.
- Lee, S., Staffileno, B. A., & Fogg, L. (2013). Influence of staff education on the function of hospitalized elders. *Nursing Outlook*, 61(1), 2-8.
- Lord, S. R., & Dayhew, J. (2001). Visual risk factors for falls in older people. *J Am Geriatr Soc*, 49(5), 508-515.
- NCCN Older Adult: Clinical Practice Guidelines. NCCN.org
- Nevitt, M. C., Cummings, S. R., Kidd, S., & Black, D. (1989). Risk factors for recurrent nonsyncopal falls. A prospective study. *JAMA*, 261(18), 2663-2668.
- Northridge, M. E., Nevitt, M. C., & Kelsey, J. L. (1996). Non-syncopal falls in the elderly in relation to home environments. *Osteoporos Int*, 6(3), 249-255.
- Overcash, J. (2007). Prediction of falls in older adults with cancer: a preliminary study. *Oncol Nurs Forum*, 34(2), 341-346. doi: 10.1188/07.onf.341-346
- Overcash, J. A., & Beckstead, J. (2008). Predicting falls in older patients using components of a comprehensive geriatric assessment. *Clin J Oncol Nurs*, 12(6), 941-949. doi: 10.1188/08.cjon.941-949

- Partridge, J. S., Fuller, M., Harari, D., Taylor, P. R., Martin, F. C., & Dhesi, J. K. (2015). Frailty and poor functional status are common in arterial vascular surgical patients and affect postoperative outcomes. *Int J Surg*, *18*, 57-63. doi: 10.1016/j.ijsu.2015.04.037
- Podsiadlo, D., & Richardson, S. (1991). The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc*, *39*(2), 142-148.
- Presley, C. J., Han, L., Leo-Summers, L., Hurria, A., Gross, C. P., Davidoff, A. J., . . . Gill, T. M. (2018). Functional trajectories before and after a new cancer diagnosis among community-dwelling older adults. *J Geriatr Oncol*. doi: 10.1016/j.jgo.2018.05.017
- Putz, M. T., Visser, M., Twisk, J. W., Deeg, D. J., & Lips, P. (2005). Endocrine and inflammatory markers as predictors of frailty. *Clin Endocrinol (Oxf)*, *63*(4), 403-411.
- Robbins, S., Waked, E., Gouw, G. J., & McClaran, J. (1994). Athletic footwear affects balance in men. *Br J Sports Med*, *28*(2), 117-122.
- Robinson, T., Walston, J., Brummel, N., Deiner, S., Brown, C., Kennedy, M., & Hurria, A., Frailty for Surgeons: Review of a National Institute on Aging Conference on Frailty for Specialties. *Journal of the American College of Surgeons*, *221* (6), 1083-1092.
- Rockwood, K., & Mitnitski, A. (2011). Frailty defined by deficit accumulation and geriatric medicine defined by frailty. *Clin Geriatr Med*, *27*(1), 17-26. doi: 10.1016/j.cger.2010.08.008
- Rubenstein, L. Z., Stuck, A. E., Siu, A. L., & Wieland, D. (1991). Impacts of geriatric evaluation and management programs on defined outcomes: overview of the evidence. *J Am Geriatr Soc*, *39*(9 Pt 2), 8S-16S; discussion 17S-18S.
- Soto-Perez-de-Celis, E., Li, D., Yuan, Y., Lau, Y. M., & Hurria, A. (2018). Functional versus chronological age: geriatric assessments to guide decision making in older patients with cancer. *Lancet Oncol*, *19*(6), e305-e316. doi: 10.1016/s1470-2045(18)30348-6
- Stenhagen, M., Ekstrom, H., Nordell, E., & Elmstahl, S. (2013). Falls in the general elderly population: a 3- and 6- year prospective study of risk factors using data from the longitudinal population study 'Good ageing in Skane'. *BMC Geriatr*, *13*, 81. doi: 10.1186/1471-2318-13-81
- Stone, C. A., Lawlor, P. G., Savva, G. M., Bennett, K., & Kenny, R. A. (2012). Prospective study of falls and risk factors for falls in adults with advanced cancer. *J Clin Oncol*, *30*(17), 2128-2133. doi: 10.1200/JCO.2011.40.7791
- Tinetti, M. E., Williams, T. F., & Mayewski, R. (1986). Fall risk index for elderly patients based on number of chronic disabilities. *Am J Med*, *80*(3), 429-434.
- Wade, D. T., & Collin, C. (1988). The Barthel ADL Index: a standard measure of physical disability? *Int Disabil Stud*, *10*(2), 64-67.
- Whitson, H. E., Sanders, L. L., Pieper, C. F., Morey, M. C., Oddone, E. Z., Gold, D. T., & Cohen, H. J. (2009). Correlation between symptoms and function in older adults with comorbidity. *J Am Geriatr Soc*, *57*(4), 676-682. doi: 10.1111/j.1532-5415.2009.02178.x

- Wildes, T. M., Maggiore, R. J., Tew, W. P., Smith, D., Sun, C. L., Cohen, H., . . . Hurria, A. (2018). Factors associated with falls in older adults with cancer: a validated model from the Cancer and Aging Research Group. *Support Care Cancer*. doi: 10.1007/s00520-018-4212-3
- Wilkins, K. (1999). Health care consequences of falls for seniors. *Health Rep*, 10(4), 47-55(ENG); 47-57(FRE).
- Wyrwich, K. W., & Wolinsky, F. D. (2000). Physical activity, disability, and the risk of hospitalization for breast cancer among older women. *J Gerontol A Biol Sci Med Sci*, 55(7), M418-421.
- Zhu, G., Zhang, X., Wang, Y., Xiong, H., Zhao, Y., & Sun, F. (2016). Effects of exercise intervention in breast cancer survivors: a meta-analysis of 33 randomized controlled trials. *Onco Targets Ther*, 9, 2153-2168. doi: 10.2147/OTT.S97864